

Recovery of Repressed Memories in Fibromyalgia Patients Treated with Hyperbaric Oxygen - Case Series Presentation and Suggested Bio-Pscho-Social Mechanism

Shai Efrati^{1*}, Amir Hadanny¹, Shir Daphna-Tekoah², Yair Bechor¹, Kobi Tiberg³, Nimrod Pik¹, Gil Suzin¹, Rachel Lev-Wiesel⁴

¹Assaf Harofeh Medical Center, Israel, ²Kaplan Medical Center, Israel, ³Loewenstein Hospital Rehabilitation Center, Israel, ⁴University of Haifa, Israel

Submitted to Journal:
Frontiers in Psychology

Specialty Section:
Movement Science and Sport Psychology

Article type:
Hypothesis and Theory Article

Manuscript ID:
325958

Received on:
30 Oct 2017

Revised on:
10 Apr 2018

Frontiers website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

SE - Study initiator, interpretation of study results, wrote the first and final drafts of the manuscript.

AH- Interpretation of study results, brain Image analysis, co-wrote with SE first and final drafts of the manuscript

SDT- Interpretation of study results, data collection, patients follow up, revision of the manuscript

YB- Data collection, patients follow up.

KT- interpretation of study results, revision of the manuscript.

NP- Patients follow up, interpretation of study results.

GS- Interpretation of study results, revision of the manuscript.

RLW- Patients follow-up, interpretation of study results, revision of the manuscript.

All authors read and approved the final manuscript.

Keywords

hyperbaric oxygen, repressed memories, Fibromyalgia, Biopsychophysical Mechanism, Childhood sexual abuse

Abstract

Word count: 245

Fibromyalgia Syndrome (FMS) is a condition considered to represent a prototype of central sensitization syndrome, characterized by chronic widespread pain and along with symptoms of fatigue, non-restorative sleep and cognitive difficulties. FMS can be induced by trauma, infection or emotional stress with cumulative evidence that dissociation is relatively frequent in FMS patients. Two randomized controlled trials have shown that hyperbaric oxygen therapy (HBOT) can induce neuroplasticity and be effectiveness in patients suffering from FMS. In this paper we present, for the first time, case series of female fibromyalgia patients who, in the course of HBOT, suddenly recalled repressed traumatic memories of childhood sexual abuse (CSA). The surfacing of the repressed memories decades after the sexual abuse events was sudden and utterly surprising. No psychological intervention was involved. As the memories surfaced, the physical pain related to FMS subsided. In one patient who had brain SPECT before and after HBOT, the prefrontal cortex appeared suppressed before and reactivated after. The 3 cases reported in this article are representative of a total of 9 fibromyalgia patients who experienced a retrieval of repressed memory during HBOT. These cases provides insights on dissociative amnesia and suggested mechanism hypothesis that is further discussed in the article. Obviously, prospective studies cannot be planned since patients are not aware of their repressed memories. However, it is very important to keep in mind the possibility of surfacing memories when treating fibromyalgia patients with HBOT or other interventions capable of awakening dormant brain regions.

Ethics statements

(Authors are required to state the ethical considerations of their study in the manuscript, including for cases where the study was exempt from ethical approval procedures)

Does the study presented in the manuscript involve human or animal subjects: Yes

Please provide the complete ethics statement for your manuscript. Note that the statement will be directly added to the manuscript file for peer-review, and should include the following information:

- Full name of the ethics committee that approved the study
- Consent procedure used for human participants or for animal owners
- Any additional considerations of the study in cases where vulnerable populations were involved, for example minors, persons with disabilities or endangered animal species

As per the Frontiers authors guidelines, you are required to use the following format for statements involving human subjects: This study was carried out in accordance with the recommendations of [name of guidelines], [name of committee]. The protocol was approved by the [name of committee]. All subjects gave written informed consent in accordance with the Declaration of

Helsinki.

For statements involving animal subjects, please use:

This study was carried out in accordance with the recommendations of 'name of guidelines, name of committee'. The protocol was approved by the 'name of committee'.

If the study was exempt from one or more of the above requirements, please provide a statement with the reason for the exemption(s).

Ensure that your statement is phrased in a complete way, with clear and concise sentences.

All reported patients, included in this article, signed informed consent and approved their case presentation.

In review

Recovery of Repressed Memories in Fibromyalgia Patients Treated with Hyperbaric Oxygen - Case Series Presentation and Suggested

Bio-Psycho-Social Biopsychophysical Mechanism

Prof. Shai Efrati, MD^{1,2,3}, Dr. Amir Hadanny, MD^{1,2}, Dr. Shir Daphna-Tekoah, PhD^{4,5}, Mr. Yair Bechor¹, Dr. Kobi Tiberg, MD⁶, Dr. Nimrod Pik, MD⁷, Mr. Gil Suzin¹, Prof. Rachel Lev-Wiesel, PhD⁸.

¹Sagol Center for Hyperbaric Medicine and Research, Assaf Harofeh Medical Center, Zerifin, Israel

²Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel

³Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

⁴Ashkelon Academic College, Ashkelon, Israel,

⁵Social Work Department, Kaplan Medical Center, Rehovot, Israel

⁶Department of Psychology, Loewenstein Center for Rehabilitation, Raanana, Israel

⁷Psychiatric Service, Assaf Harofeh Medical Center, Zerifin, Israel.

⁸The Emili Sagol CAT Research Center, Graduate School of Creative Arts Therapies University of Haifa

Running head: Repressed memories recovered during hyperbaric oxygen therapy

Word count: 3634

Corresponding author:

Professor Shai Efrati, MD

Head of the Sagol Center for Hyperbaric Medicine and Research,

Head of Research & Development,

Assaf-Harofeh Medical Center, Israel

Associate professor, Sackler School of Medicine and Sagol School of Neuroscience,

Tel-Aviv University, Tel-Aviv, Israel

Mail: efratishai@013.net

Phone: +972-(0)8-977-9393/5. FAX: +972-(0)8-9542238

Abstract

Fibromyalgia Syndrome (FMS) is a condition considered to represent a prototype of central sensitization syndrome, characterized by chronic widespread pain and along with symptoms of fatigue, non-restorative sleep and cognitive difficulties. FMS can be induced by trauma, infection or emotional stress with cumulative evidence that dissociation is relatively frequent in FMS patients. Two randomized controlled trials have shown that hyperbaric oxygen therapy (HBOT) can induce neuroplasticity and be effectiveness in patients suffering from FMS. In this paper we present, for the first time, case series of female fibromyalgia patients who, in the course of HBOT, suddenly recalled repressed traumatic memories of childhood sexual abuse (CSA). The surfacing of the repressed [\(dissociative\)](#) memories decades after the sexual abuse events was sudden and utterly surprising. No psychological intervention was involved. As the memories surfaced, the physical pain related to FMS subsided. In one patient who had brain SPECT before and after HBOT, the prefrontal cortex appeared suppressed before and reactivated after. The 3 cases reported in this article are representative of a total of 9 fibromyalgia patients who experienced a retrieval of repressed memory during HBOT. These cases provides insights on dissociative amnesia and suggested mechanism hypothesis that is further discussed in the article. Obviously, prospective studies cannot be planned since patients are not aware of their repressed memories. However, it is very important to keep in mind the possibility of surfacing memories when treating fibromyalgia patients with HBOT or other interventions capable of awakening dormant brain regions.

Key words: Childhood sexual abuse, Repressed memories, Fibromyalgia, Hyperbaric oxygen

Introduction

Fibromyalgia Syndrome (FMS) is a condition characterized by chronic widespread pain and diffuse tenderness, along with symptoms of fatigue, non-restorative sleep and cognitive difficulties. It affects 2-45% of the [general population worldwide](#), with 9:1 female-to-male incidence ratio (Buskila 2009, Branco, Bannwarth et al. 2010, Clauw, Arnold et al. 2011, Schmidt-Wilcke and Clauw 2011, Jones, Atzeni et al. 2015). FMS is considered to represent a prototype of central sensitization, i.e. a condition characterized by an increase in the transmission and processing of pain within the central nervous system (Yunus 2007, Yunus 2007, Hadanny and Efrati 2015, Ablin, Efrati et al. 2016). Like other functional somatic syndromes, despite suffering from pain affecting soft tissues (muscles, ligaments), and tendons, patients seem well without obvious abnormalities on neither physical examination nor evidence of tissue inflammation on laboratory and radiologic studies (Clauw 2014).

In recent years, there are cumulative data regarding the complex biology of chronic pain within the CNS (Ablin, Efrati et al. 2016, Sancassiani, Machado et al. 2017, Yavne, Amital et al. 2018). Chronic pain syndromes in general, and FMS in particular, are not congenital; these conditions rather evolve over lifetime, often in response to various external factors such as physical trauma, infection or emotional stress, which emphasizes the capability of the CNS to morph and re-wire during life, even at a fully developed stage (Ablin, Efrati et al. 2016, Yavne, Amital et al. 2018). Hence, the general concept of neuro-plasticity is an essential pattern of chronic pain evolution.

In this study, our main focus is on FMS induced by childhood sexual abuse (CSA). CSA has negative long term physical and psychological effects and women with a history of CSA may develop depression, post-traumatic stress disorder (PTSD),

dissociative disorders and chronic pain syndromes such as FMS (Spiegel, Chatterjee et al. 2015). Dissociation in CSA is particularly relevant to amnesia of all or part of the traumatic events (Bohn, Bernardy et al. 2013). Studies have shown that dissociation is relatively frequent in FMS patients (Naring, van Lankveld et al. 2007, Bohn, Bernardy et al. 2013, Kilic, Sar et al. 2014, Berkol, Balcioglu et al. 2017). Unfortunately, despite the fact that chronic pain syndromes are common in adult survivors of CSA, patients rarely reveal their abuse history to their care providers due to amnesia, dissociation, or merely shame, embarrassment or trust issues.

HBOT, the application of hyperbaric pressure in conjunction with increased oxygen content, has been shown in several clinical studies to have the capacity to induce neuroplasticity in injured brains even years after an acute insult (Boussi-Gross, Golan et al. 2013, Efrati, Fishlev et al. 2013, Efrati and Ben-Jacob 2014, Boussi-Gross, Golan et al. 2015, Hadanny, Golan et al. 2015, Tal, Hadanny et al. 2015, Hadanny and Efrati 2016). As demonstrated in different animal models, HBOT induces neuroplasticity by stimulating cell proliferation (Mu, Ostrowski et al. 2013), promotes neurogenesis of endogenous neural stem cells (Yang, Wang et al. 2008), regenerates axonal white matter (Chang, Lee et al. 2009), improves maturation and myelination of injured peripheral and cranial neural fibers (Haapaniemi, Nylander et al. 1998, Vilela, Lazarini et al. 2008), induces brain angiogenesis (Tal, Hadanny et al. 2015), and stimulates axonal growth (Mukoyama, Iida et al. 1975, Bradshaw, Nelson et al. 1996).

Despite the fact that HBOT for neurological disorders is somewhat still controversial there is recent evidence of its effectiveness in treating FMS (Ablin, Efrati et al. 2016). To date, two prospective randomized controlled trials have demonstrated the efficacy of HBOT in fibromyalgia (Yildiz, Kiralp et al. 2004, Hadanny, Golan et al. 2015). The

improvement was demonstrated in all aspects of FMS including pain threshold, fatigue, distress and quality of life (Yildiz, Kiralp et al. 2004, Hadanny, Golan et al. 2015).

In this paper we present three cases of women suffering from fibromyalgia, who recalled suppressed traumatic memories of CSA during HBOT. These cases provides additional important insights on memories retrieval and dissociative amnesia and the opportunity to suggest a possible mechanism hypothesis that is further discussed in the article. The participants in the current paper were not subject to any psychological intervention during the time of HBOT. As the memories surfaced, around the 20th HBOT session (with more memories surfacing as the treatment continued) the physical pain related to the FMS subsided. Following participants' disclosure to a nurse, they were assigned to a physician for further assistance and investigation.

It should be noted that these events of self-disclosure were unfamiliar and unexpected for both patients and medical staff.

Material and Methods

A retrospective case presentation of FMS patients who reported retrieval of repressed memories during HBOT at the Sagol center for hyperbaric medicine and research at Assaf-Harofeh medical center, Israel. The three cases reported in this article represent 9 FMS patients who experienced the same during HBOT, out of a larger cohort of ~200 FMS patients treated by HBOT in our center between 2013-2016. All reported patients, included in this article, signed [an informed consent for their case publication](#) [After patients signed consent, they reviewed and proofed their presented case. There were six additional patients with repressed memories recovery who did not agree for case publication. Thus, their cases are not presented. Vulnerable populations \(for example minors, persons with disabilities\) were not involved in this study. The publication](#)

process of those cases was approved by the head of Assaf Harofeh Medical Center Institutional Review board. and approved their case presentation. The additional 6 cases did not agree their case would be published and thus not presented.

The following HBOT protocol was practiced: 60 daily sessions, 5 days a week, 90 minutes each session, breathing 100% oxygen at 2.0 absolute atmospheres (ATA) with 5 minutes air breaks every 30 minutes. Sessions were performed in "multiplace" chambers, which may treat up to 12 patients simultaneously.

The diagnosis of FMS was based on the following criteria: (1) Symptoms of widespread pain, both above and below the waist and bilaterally. (2) Physical findings of at least 11 out of a set of 18 known tender points.

Results

Case study 1

N, a 21 year-old female, was diagnosed at the age of seventeen with severe FMS, because of which she had to quit studying. N was a popular, good student, a third-born of three children to her highly educated parents. At the age of fourteen she insisted on moving to another school, rationalizing it as her simply wishing to study at a "better school". Her parents agreed. She then invested all her energy in studying and in dancing lessons, in which she excelled until the onset of her fibromyalgia. N was referred to the Sagol hyperbaric institute by the family physician to receive HBOT for her fibromyalgia. Prior to her referral to HBOT she was treated with several different medications and with medical Cannabis for her severe pain. Between the 17 and the 34 HBOT sessions, N complained on low energy and tiredness. In additional, N told the medical staff that she feels sadder and more anxious than prior to HBOT. During that

time, she started to experience flashbacks of herself being sexually molested. The flashbacks occurred both during HBOT sessions (inside the chamber) and in the hours following a HBOT session (at her normal environment). The flashbacks started with unbearable sexual arousal, accompanied with shortness of breath, burst into tears and other panic attack related symptoms such as feeling of loss of control and overwhelming fear. Fragments of memories appeared, in which she was being raped by several peers which added up to a full recovery of her repressed memory of been repeatedly molested and raped by classmates at the age of fourteen.

Looking back, N said: "I can't understand how I, or anyone else for that matter, could forget such experiences... I continued to function as an excellent student; I focused on my dancing... I avoided peer gatherings, rationalizing this avoidance to myself and to my parents as "lack of interest" and "will to excel."

N reported that, after the repressed memories emerged, she felt much stronger, slept better, had more energy and had almost no pain. According to N, she was fully recovered from her FMS.

Case study 2

A, a 56 year-old businesswoman, married with four children, the eldest daughter in her family of origin. As a child she had urolithiasis, for which she underwent bladder surgery three times. The relationships between her parents were complicated. Her mother used to have love affairs and brought her sexual partners home in the father's absence. The mother also used to share her sexual experiences with A, who was a "parental child". At the age of seven, A has been visiting an eighteen year-old neighbor in his apartment three times a week and was receiving little presents from him. After the first visit, A cut her long, beautiful, blond hair without informing her parents, telling

them later that she "preferred to be ugly". About a year later, during the Six Days War, the family moved for a week to her grandmother's. A describes this week as "the best, quietest, calmest week of my life, despite the fear of being at war." Upon returning home, she refused her neighbor's invitations and forgot his name. Shortly after graduation she got married. In 2011, she had a car accident and referred to treatment at the Sagol HBOT institute. Around that time she was also diagnosed with FMS. Between the 37th and 44th HBOT sessions, A experienced flashbacks of sexual molestation. The flashbacks were accompanied with anxiety and feelings of losing control in addition to sympathetic signs such as sweating. The flashbacks began with unbearable sexual arousal and concurrent panic attacks. Flashbacks initially occurred during the hyperbaric sessions, and later on happened during her daily routines between sessions. Visually vivid memories surfaced including her being ordered to have oral intercourse. These flashbacks gradually joined into detailed, coherent memories of sexual molestation she had experienced at the age of seven by her eighteen year-old neighbor, who threatened to kill her if she ever revealed their "secret". In an attempt to validate these memories, A returned to her childhood neighborhood, found the perpetrator, and confronted him. He confirmed that this had indeed occurred as remembered.

Case study 3

B, a 36 year-old female, was the youngest of three siblings in her family of origin. When admitted to HBOT for fibromyalgia, she reported having been sexually abused by her uncle (her mother's brother) between the ages of six and sixteen. She said that her parents didn't know about the abuse. "Actually", she said "I think that, back then, I myself did not remember it most of the day, kind of detached." Her mother died of cancer a year after the abuse ceased. B described her mother as "an innocent, good

mother who knew nothing of what was going on." In adulthood, B had psychotherapy for her CSA. During her Social Work studies, B began to suffer from fibromyalgia. The disease prevented her from committing to a full-time work as a social worker. At the age of thirty-six, she started HBOT for her FMS. During the first twenty sessions, she reported sexual arousal and unbearable pain in her abdomen, "felt as if someone was stabbing me with a knife". These symptoms were followed by surfacing of new memories of the abuse revealing another abuser - her mother. "My mother took me to the shower after he [the uncle] abused me. She undressed me, ordered me to open my legs and then began to abuse my genitals." These new, vivid, coherent and detailed memories exposed her to different narratives on her mother and her childhood. Those memories of abuse by B's mother did not surface prior to the HBOT, even during psychotherapy. After those new memories surfaced she returned to psychotherapy.

B's baseline and post HBOT brain SPECT scans are presented in Figure 1. The highest relative increases in activity (>15%) are seen in the visual cortex (BA 19), prefrontal cortex (BA 9), orbito-frontal gyrus (BA 11) and paratemporal cortex (BA37) (Figure-1).

Discussion

This article is the first report of retrieval of repressed memories of CSA in FMS patients during HBOT. In the three cases presented, HBOT was initiated for FMS, decades after the sexual abuse events occurred. The retrieval of the repressed memory through physiological intervention that induces neuroplasticity was unexpected. This phenomenon may augment our knowledge on the pathophysiological process related to

dissociation and dissociative amnesia and suggest new physiological interventions to enhance the current psychological interventions.

Dissociation, Dissociative amnesia and Biopsychophysical mechanism

Trauma is a biopsychophysical experience, even when the traumatic event doesn't cause direct mechanical harm to the body. According to this, the organism is defined by a collection of multiple biological and psychological interactions as well as social factors throughout its life (Bowman 2011, Nijenhuis and van der Hart 2011). The term "Dissociation" refers to the failure of integrating aspects of memory, identity, perception, and consciousness. Dissociative amnesia, is a failure of the ability to remember autobiographical information in the absence of brain damage (Staniloiu and Markowitsch 2014). Thus "Dissociation" means that a memory is neither lost nor forgotten but is unavailable for retrieval for a period of time, which could last even decades (Chu, Frey et al. 1999). Dissociative amnesia or the suppression of traumatic memories, is well-documented in the literature and is coded as a dissociative disorder in Diagnostic and Statistical Manual of Mental Disorders (DSM-5®), Fifth Edition (American Psychiatric Association. and American Psychiatric Association. DSM-5 Task Force. 2013).

Individuals with dissociative disorders usually report a history of exposure to traumatic stress, development trauma, or other events that trigger stress known as stressors (Andrews, Brewin et al. 1999, Bohn, Bernardy et al. 2013, Fergusson, McLeod et al. 2013, Lev-Wiesel 2014). In psychological terms, dissociation may be considered as a protective mechanism to shield the traumatized individual from the unbearable pain of the memory and its consequences. CSA is an exceptionally severe form of emotional and physical trauma that may lead to disorders of memory and dissociation (Andrews,

Brewin et al. 1999, McNally 2003, van der Kolk, Roth et al. 2005, Fergusson, McLeod et al. 2013, Lev-Wiesel 2014). Several clinical studies demonstrate that CSA is associated with FMS severity and may shape the biological development of interception in ways that predispose to pain and polysymptomatic distress (Romans, Belaise et al. 2002, Ciccone, Elliott et al. 2005, Jiao, Vincent et al. 2015, Ortiz, Ballard et al. 2016). The classic fight-or-flight response to a perceived threat is a reflexive neural phenomenon which has obvious survival advantages in evolutionary terms (Sherin and Nemeroff 2011). However, under certain circumstances, such as the CSA cases described above, exposure to a perceived threat can cause dysregulated response (Scaer 2001, Sherin and Nemeroff 2011). Chronic dysregulation can lead to functional impairment in certain individuals who become “psychologically traumatized” with long standing neurobiological abnormalities, which overlap features found in patients with post-concussion syndrome due to traumatic brain injury (TBI) (Scaer 2001, Sherin and Nemeroff 2011). Certain brain areas become disconnected from the normal trophic stimulations of cerebral perception and subjected to the extremes of vasomotor instability of trauma that may lead to pathologic vasoconstriction and ischemia (Scaer 2001). This model explains the somatoform" dissociative symptomatology that may arise in TBI or CSA, followed by FMS. Thus, FMS can be induced by either TBI or CSA.

Suggested physiological mechanism for HBOT effect in dissociation

From the neurobiological/physiological perspective, the authors suggest a possible mechanism for the recovery of repressed memories induced by HBOT.

Astrup et al. demonstrated in 1977, in a paper argued to be one of the most important in applied neurology, that after the onset of focal ischemia (such as blood vessel

occlusion), measurements of electrical activity reveal brain regions that are dysfunctional but still viable (Astrup, Symon et al. 1977). The neurons in this area retained enough energy to maintain ion pumps and sustain the -70-mV resting membrane potentials required for their existence. However, those neurons did not have enough energy to generate the action potentials needed for their purpose functioning. In the last decade, functional imaging of the brain by Positron Emission Tomography (PET) and Single Photon Emission CT (SPECT) scanning (Meerwaldt, Slart et al. , Heiss 2000, Baron 2001, Siddique, Fernandes et al. 2002, Ebinger, De Silva et al. 2009), afforded us images of such areas, and the accumulating data indicate that apparently dead areas may persist for years after an acute event (Boussi-Gross, Golan et al. 2013, Efrati, Fishlev et al. 2013). The stunned/hibernating areas, unable to fire action potentials, is characterized by metabolic dysfunction, namely anaerobic metabolism and ATP depletion (Figure-2). Loss of ATP and the resultant intracellular acidosis would bring about energetic breakdown of ion pumps and, consequently, additional damage to the mitochondria and endoplasmic reticulum (Hossmann 2006, Culmsee and Krieglstein 2007). Furthermore, anaerobic metabolism would bring about elevation of free radical levels (Hossmann 2006, Culmsee and Krieglstein 2007), persistent inhibition of protein synthesis (DeGracia 2004), selective neuronal damage demonstrable by decreased density of benzodiazepine and/or 5-HT₂ receptors (Vera, Zilbovicius et al. 1996, Yamauchi, Kudoh et al. 2005), augmented intracellular calcium (Hossmann 2006), blood-brain barrier damage and augmented inflammation evidenced by higher levels of cytokines and cyclooxygenase-2 (COX-2) (Yin, Badr et al. 2002). The brain regions with continuous hypoperfusion, with their very low, anaerobic metabolic rate, settle into a quasi-steady energy well from which they cannot climb up on their own. The hibernation state typical of some mammalian species seems to be the

most proper model created by nature, simulating the post injury and post ischemia states of brain (Drew, Buck et al. 2007). The energy consumption in hibernation is less than 0.1 WT/kg, i.e. less than 1% of the metabolism in a resting state. The corresponding outcomes at the cellular level include a dramatic (~2500-fold) decrease in protein synthesis (Frerichs, Smith et al. 1998), complete interruption of neuronal spike activity (Carey, Andrews et al. 2003), and a switch from carbohydrate-based to fat-based metabolism (Carey, Andrews et al. 2003). In turn, metabolic changes induced by minimized levels of energy availability have an impact on brain structure and functioning at the cellular level, such as reorganization of cell membranes, formation of protein-free domains that displace membrane proteins, and altered cell membrane permeability due to alterations in cytoplasmic matrix (Azzam, Hallenbeck et al. 2000, Adibhatla and Hatcher 2008).

The neurobiological mechanism of dissociation, as mentioned above involves disconnection of certain brain areas, may be explained by hibernation, a hypometabolic state at the cellular level (minimal energy generation by the mitochondria in order to maintain membrane potential for cell survival). Hibernated areas may correlate with apathy and low responsiveness (Schore 2009). This model supports the assumption that dissociation could be interpreted from the biological perspective as a hypometabolic state and from the biopsychological perspective as a deficiency of psychological energy. In the cases presented, the metabolic dysfunction of the dissociated brain regions serves as a physiological boundary for psychological intervention (Figure-2).

Energy wise, climbing back up from the "metabolic well" mandates energy input. As discussed above, HBOT has been shown to induce neuroplasticity and reactivation of cells in chronic metabolic dysfunction in different type of brain injuries, and in FMS specifically (Mukoyama, Iida et al. 1975, Bradshaw, Nelson et al. 1996, Haapaniemi,

Nylander et al. 1998, Yildiz, Kiralp et al. 2004, Vilela, Lazarini et al. 2008, Yang, Wang et al. 2008, Chang, Lee et al. 2009, Boussi-Gross, Golan et al. 2013, Efrati, Fishlev et al. 2013, Mu, Ostrowski et al. 2013, Efrati and Ben-Jacob 2014, Boussi-Gross, Golan et al. 2015, Hadanny and Efrati 2015, Hadanny, Golan et al. 2015, Tal, Hadanny et al. 2015, Hadanny and Efrati 2016). HBOT supplies extra oxygen to the anaerobic brain regions, providing them with the energy needed to exit metabolic well. Once these regions of metabolic dysfunction are re-activated by HBOT, cerebral blood flow is increased to these regions and they return to normal metabolism. Hibernating brain regions which may have been responsible to memory repression, can be reactivated by HBOT, regain their normal function and resurface suppressed memories (Figure-3).

Potential role of prefrontal cortex role in suppressed memories

One of the patients (case 1) presented in this article had brain SPECT before and after HBOT (Figure1). In this case, the prefrontal cortex that was suppressed at baseline was reactivated by the use of HBOT.

Repression of memories, or dissociative amnesia, has been correlated with decreased activity of the dorsomedial prefrontal cortex, while recovery is correlated with increased prefrontal cortex activity as shown by functional magnetic resonance imaging (fMRI) (Anderson, Ochsner et al. 2004, Kuhl, Kahn et al. 2008, Che, Luo et al. 2015). PET studies in patients suffering from PTSD demonstrated increased cerebral blood flow in the prefrontal cortex when re-experiencing emotionally charged episodic memories (YOSHIHIRO MASAKI 2006, Im, Namgung et al. 2016).

Study limitations

This article has several obvious limitations. Most limitations are related to the small number of patients and the lack of brain SPECT imaging in the other two patients. Another limitation relates to the fact that there is no known way to fully and objectively validate the retrieved memories. Still, two of the women confronted their past perpetrators, who begged for their forgiveness. The three cases presented in this article are representative of 9 women who had retrieved suppressed memories during HBOT at our center. In order to gain better understanding of dissociation/dissociative amnesia and its suggested biopsychophysical mechanism and long term biological consequences, direct evaluation of brain metabolism/activity in animal models should be investigated.

To conclude, this is the first clinical report of recovery of suppressed memories with HBOT alone as physiological intervention. Even though further studies are needed, it would not be easy to carry out a study in patients that are unaware of their difficult childhood history. Obviously, prospective studies cannot be planned since patients are not aware of their repressed memories. However, it is very important to keep in mind the possibility of surfacing memories when treating FMS patients with HBOT or other interventions capable of awakening dormant brain regions. We recommend that HBOT practitioners treating FMS patients should add the potential risk for unexpected memory of traumatic events to occur to their informed consent. All medical professionals who are dealing with FMS patients should be aware and prepared to handle such outcome. It would be highly beneficial if anyone who encounters such a case documents and reports it. Through the cumulative data we may get important insight into the neurobiology of suppressed memories.

Acknowledgement statement: We are thankful Dr. Ilan Kutz for enlightening discussions regarding the neurophysiology effects of HBOT in this unique condition. We are very thankful for Michal Ben-Jacob for her critical reading and editing the two versions of the manuscript.

In review

Figure-1: Case 1 SPECT calculated change after HBOT compared to baseline. In the top two rows colors, represent maximal functional brain activity relative to the entire brain activity; in the bottom row, colors represent regional changes in functional brain activity. White and red areas show the biggest relative changes in cerebral blood flow/perfusion. Significant changes can be seen at the right prefrontal and temporal areas (white).

Abbreviations:

Analysis of brain SPECT performed pre- and post HBOT showed the highest relative increase in activity (>15%) in the visual cortex (BA 19), prefrontal cortex (BA 9), orbito-frontal gyrus (BA 11) and paratemporal cortex (BA37).

Figure 2: Dissociation and repressed memory - suggested model for the neuropathology cascade.

Figure-3: Recovery of repressed memory by HBOT – suggested mechanism

References

- Ablin, J. N., S. Efrati and D. Buskila (2016). "Building up the pressure on chronic pain." Clinical and experimental rheumatology **34**(2 Suppl 96): S3-5.
- Ablin, J. N., S. Efrati and D. Buskila (2016). "Building up the pressure on chronic pain." Clin Exp Rheumatol **34**(2 Suppl 96): S3-5.
- Adibhatla, R. M. and J. F. Hatcher (2008). "Altered lipid metabolism in brain injury and disorders." Subcell Biochem **49**: 241-268.
- American Psychiatric Association. and American Psychiatric Association. DSM-5 Task Force. (2013). Diagnostic and statistical manual of mental disorders : DSM-5. Washington, D.C., American Psychiatric Publishing.
- Anderson, M. C., K. N. Ochsner, B. Kuhl, J. Cooper, E. Robertson, S. W. Gabrieli, G. H. Glover and J. D. Gabrieli (2004). "Neural systems underlying the suppression of unwanted memories." Science **303**(5655): 232-235.
- Andrews, B., C. R. Brewin, J. Ochera, J. Morton, D. A. Bekerian, G. M. Davies and P. Mollon (1999). "Characteristics, context and consequences of memory recovery among adults in therapy." The British journal of psychiatry : the journal of mental science **175**: 141-146.
- Astrup, J., L. Symon, N. M. Branston and N. A. Lassen (1977). "Cortical evoked potential and extracellular K⁺ and H⁺ at critical levels of brain ischemia." Stroke **8**(1): 51-57.
- Azzam, N. A., J. M. Hallenbeck and B. Kachar (2000). "Membrane changes during hibernation." Nature **407**(6802): 317-318.
- Baron, J. C. (2001). "Mapping the ischaemic penumbra with PET: a new approach." Brain **124**(Pt 1): 2-4.
- Berkol, T. D., Y. H. Balcioglu, S. S. Kirlioglu, H. Erensoy and M. Vural (2017). "Dissociative features of fibromyalgia syndrome." Neurosciences (Riyadh) **22**(3): 198-204.
- Bohn, D., K. Bernardy, F. Wolfe and W. Hauser (2013). "The association among childhood maltreatment, somatic symptom intensity, depression, and somatoform dissociative symptoms in patients with fibromyalgia syndrome: a single-center cohort study." J Trauma Dissociation **14**(3): 342-358.
- Boussi-Gross, R., H. Golan, G. Fishlev, Y. Bechor, O. Volkov, J. Bergan, M. Friedman, D. Hoofien, N. Shlamkovitch, E. Ben-Jacob and S. Efrati (2013). "Hyperbaric oxygen therapy can improve post concussion syndrome years after mild traumatic brain injury - randomized prospective trial." PLoS One **8**(11): e79995.
- Boussi-Gross, R., H. Golan, O. Volkov, Y. Bechor, D. Hoofien, M. S. Beerli, E. Ben-Jacob and S. Efrati (2015). "Improvement of memory impairments in poststroke patients by hyperbaric oxygen therapy." Neuropsychology **29**(4): 610-621.
- Bowman, E. S. (2011). "Nijenhuis and Van der Hart: one view of the elephant." J Trauma Dissociation **12**(4): 446-449; discussion 469-473.
- Bradshaw, P. O., A. G. Nelson, J. W. Fanton, T. Yates and K. S. Kagan-Hallet (1996). "Effect of hyperbaric oxygenation on peripheral nerve regeneration in adult male rabbits." Undersea Hyperb Med **23**(2): 107-113.
- Branco, J. C., B. Bannwarth, I. Failde, J. Abello Carbonell, F. Blotman, M. Spaeth, F. Saraiva, F. Nacci, E. Thomas, J. P. Caubere, K. Le Lay, C. Taieb and M. Matucci-Cerinic (2010). "Prevalence of fibromyalgia: a survey in five European countries." Semin Arthritis Rheum **39**(6): 448-453.
- Buskila, D. (2009). "Developments in the scientific and clinical understanding of fibromyalgia." Arthritis research & therapy **11**(5): 242.

Carey, H. V., M. T. Andrews and S. L. Martin (2003). "Mammalian hibernation: cellular and molecular responses to depressed metabolism and low temperature." Physiol Rev **83**(4): 1153-1181.

Chang, C. C., Y. C. Lee, W. N. Chang, S. S. Chen, C. C. Lui, H. W. Chang, W. L. Liu and Y. L. Wang (2009). "Damage of white matter tract correlated with neuropsychological deficits in carbon monoxide intoxication after hyperbaric oxygen therapy." J Neurotrauma **26**(8): 1263-1270.

Che, X., X. Luo, D. Tong, B. M. Fitzgibbon and J. Yang (2015). "Habitual suppression relates to difficulty in regulating emotion with cognitive reappraisal." Biol Psychol **112**: 20-26.

Chu, J. A., L. M. Frey, B. L. Ganzel and J. A. Matthews (1999). "Memories of childhood abuse: dissociation, amnesia, and corroboration." The American journal of psychiatry **156**(5): 749-755.

Ciccione, D. S., D. K. Elliott, H. K. Chandler, S. Nayak and K. G. Raphael (2005). "Sexual and physical abuse in women with fibromyalgia syndrome: a test of the trauma hypothesis." The Clinical journal of pain **21**(5): 378-386.

Clauw, D. J. (2014). "Fibromyalgia: a clinical review." JAMA **311**(15): 1547-1555.

Clauw, D. J., L. M. Arnold and B. H. McCarberg (2011). "The science of fibromyalgia." Mayo Clinic proceedings **86**(9): 907-911.

Culmsee, C. and J. Kriegelstein (2007). "Ischaemic brain damage after stroke: new insights into efficient therapeutic strategies. International Symposium on Neurodegeneration and Neuroprotection." EMBO Rep **8**(2): 129-133.

DeGracia, D. J. (2004). "Acute and persistent protein synthesis inhibition following cerebral reperfusion." J Neurosci Res **77**(6): 771-776.

Drew, K. L., C. L. Buck, B. M. Barnes, S. L. Christian, B. T. Rasley and M. B. Harris (2007). "Central nervous system regulation of mammalian hibernation: implications for metabolic suppression and ischemia tolerance." J Neurochem **102**(6): 1713-1726.

Ebinger, M., D. A. De Silva, S. Christensen, M. W. Parsons, R. Markus, G. A. Donnan and S. M. Davis (2009). "Imaging the penumbra - strategies to detect tissue at risk after ischemic stroke." J Clin Neurosci **16**(2): 178-187.

Efrati, S. and E. Ben-Jacob (2014). "Reflections on the neurotherapeutic effects of hyperbaric oxygen." Expert review of neurotherapeutics **14**(3): 233-236.

Efrati, S., G. Fishlev, Y. Bechor, O. Volkov, J. Bergan, K. Kliakhandler, I. Kamiager, N. Gal, M. Friedman, E. Ben-Jacob and H. Golan (2013). "Hyperbaric oxygen induces late neuroplasticity in post stroke patients--randomized, prospective trial." PLoS One **8**(1): e53716.

Fergusson, D. M., G. F. McLeod and L. J. Horwood (2013). "Childhood sexual abuse and adult developmental outcomes: findings from a 30-year longitudinal study in New Zealand." Child abuse & neglect **37**(9): 664-674.

Frerichs, K. U., C. B. Smith, M. Brenner, D. J. DeGracia, G. S. Krause, L. Marrone, T. E. Dever and J. M. Hallenbeck (1998). "Suppression of protein synthesis in brain during hibernation involves inhibition of protein initiation and elongation." Proc Natl Acad Sci U S A **95**(24): 14511-14516.

Haapaniemi, T., G. Nylander, M. Kanje and L. Dahlin (1998). "Hyperbaric oxygen treatment enhances regeneration of the rat sciatic nerve." Exp Neurol **149**(2): 433-438.

Hadanny, A. and S. Efrati (2015). "Oxygen--a limiting factor for brain recovery." Crit Care **19**: 307.

Hadanny, A. and S. Efrati (2016). "Treatment of persistent post-concussion syndrome due to mild traumatic brain injury: current status and future directions." Expert review of neurotherapeutics **16**(8): 875-887.

Hadanny, A., H. Golan, G. Fishlev, Y. Bechor, O. Volkov, G. Suzin, E. Ben-Jacob and S. Efrati (2015). "Hyperbaric oxygen can induce neuroplasticity and improve cognitive functions of

patients suffering from anoxic brain damage." Restorative neurology and neuroscience **33**(4): 471-486.

Hadanny, A., H. Golan, G. Fishlev, Y. Bechor, O. Volkov, G. Suzin, E. Ben-Jacob and S. Efrati (2015). "Hyperbaric oxygen can induce neuroplasticity and improve cognitive functions of patients suffering from anoxic brain damage." Restor Neurol Neurosci **33**(4): 471-486.

Heiss, W. D. (2000). "Ischemic penumbra: evidence from functional imaging in man." J Cereb Blood Flow Metab **20**(9): 1276-1293.

Hossmann, K. A. (2006). "Pathophysiology and therapy of experimental stroke." Cell Mol Neurobiol **26**(7-8): 1057-1083.

Im, J. J., E. Namgung, Y. Choi, J. Y. Kim, S. J. Rhie and S. Yoon (2016). "Molecular Neuroimaging in Posttraumatic Stress Disorder." Experimental neurobiology **25**(6): 277-295.

Jiao, J., A. Vincent, S. S. Cha, C. A. Luedtke and T. H. Oh (2015). "Association of abuse history with symptom severity and quality of life in patients with fibromyalgia." Rheumatology international **35**(3): 547-553.

Jones, G. T., F. Atzeni, M. Beasley, E. Fluss, P. Sarzi-Puttini and G. J. Macfarlane (2015). "The prevalence of fibromyalgia in the general population: a comparison of the American College of Rheumatology 1990, 2010, and modified 2010 classification criteria." Arthritis Rheumatol **67**(2): 568-575.

Kilic, O., V. Sar, O. Taycan, C. Aksoy-Poyraz, T. C. Erol, O. Tecer, M. H. Emul and M. Ozmen (2014). "Dissociative depression among women with fibromyalgia or rheumatoid arthritis." J Trauma Dissociation **15**(3): 285-302.

Kuhl, B. A., I. Kahn, N. M. Dudukovic and A. D. Wagner (2008). "Overcoming suppression in order to remember: contributions from anterior cingulate and ventrolateral prefrontal cortex." Cogn Affect Behav Neurosci **8**(2): 211-221.

Lev-Wiesel, R., Amir G. (2014). "The Role of Dissociation in Self-Injurious Behavior among Female Adolescents Who Were Sexually Abused." Journal of Child Sexual Abuse **23**(7): 18

McNally, R. J. (2003). "Psychological mechanisms in acute response to trauma." Biol Psychiatry **53**(9): 779-788.

Meerwaldt, R., R. H. Slart, G. M. van Dam, G. J. Luijckx, R. A. Tio and C. J. Zeebregts "PET/SPECT imaging: from carotid vulnerability to brain viability." Eur J Radiol **74**(1): 104-109.

Mu, J., R. P. Ostrowski, Y. Soejima, W. B. Rolland, P. R. Krafft, J. Tang and J. H. Zhang (2013). "Delayed hyperbaric oxygen therapy induces cell proliferation through stabilization of cAMP responsive element binding protein in the rat model of MCAo-induced ischemic brain injury." Neurobiol Dis **51**: 133-143.

Mukoyama, M., M. Iida and I. Sobue (1975). "Hyperbaric oxygen therapy for peripheral nerve damage induced in rabbits with cloquinol." Exp Neurol **47**(3): 371-380.

Naring, G. W., W. van Lankveld and R. Geenen (2007). "Somatoform dissociation and traumatic experiences in patients with rheumatoid arthritis and fibromyalgia." Clin Exp Rheumatol **25**(6): 872-877.

Nijenhuis, E. R. and O. van der Hart (2011). "Dissociation in trauma: a new definition and comparison with previous formulations." J Trauma Dissociation **12**(4): 416-445.

Ortiz, R., E. D. Ballard, R. Machado-Vieira, L. N. Saligan and B. Walitt (2016). "Quantifying the influence of child abuse history on the cardinal symptoms of fibromyalgia." Clinical and experimental rheumatology **34**(2 Suppl 96): S59-66.

Romans, S., C. Belaise, J. Martin, E. Morris and A. Raffi (2002). "Childhood abuse and later medical disorders in women. An epidemiological study." Psychotherapy and psychosomatics **71**(3): 141-150.

Sancassiani, F., S. Machado, V. Ruggiero, E. Cacace, C. Carmassi, C. Gesi, L. Dell'Osso and M. G. Carta (2017). "The management of fibromyalgia from a psychosomatic perspective: an overview." Int Rev Psychiatry **29**(5): 473-488.

Scaer, R. C. (2001). "The neurophysiology of dissociation and chronic disease." Appl Psychophysiol Biofeedback **26**(1): 73-91.

Schmidt-Wilcke, T. and D. J. Clauw (2011). "Fibromyalgia: from pathophysiology to therapy." Nature reviews. Rheumatology **7**(9): 518-527.

Schore, A. N. (2009). Attachment trauma and the developing right brain: Origins of pathological dissociation. Dissociation and the dissociative disorders: DSM-V and beyond J. A. O. N. P.F. Dell. New York, Routledge: 107-141.

Sherin, J. E. and C. B. Nemeroff (2011). "Post-traumatic stress disorder: the neurobiological impact of psychological trauma." Dialogues Clin Neurosci **13**(3): 263-278.

Siddique, M. S., H. M. Fernandes, T. D. Wooldridge, J. D. Fenwick, P. Slomka and A. D. Mendelow (2002). "Reversible ischemia around intracerebral hemorrhage: a single-photon emission computerized tomography study." J Neurosurg **96**(4): 736-741.

Spiegel, D. R., A. Chatterjee, A. L. McCroskey, T. Ahmadi, D. Simmelink, E. C. t. Oldfield, C. R. Pryor, M. Faschan and O. Raulli (2015). "A Review of Select Centralized Pain Syndromes: Relationship With Childhood Sexual Abuse, Opiate Prescribing, and Treatment Implications for the Primary Care Physician." Health Serv Res Manag Epidemiol **2**: 2333392814567920.

Staniloiu, A. and H. J. Markowitsch (2014). "Dissociative amnesia." Lancet Psychiatry **1**(3): 226-241.

Tal, S., A. Hadanny, N. Berkovitz, E. Sasson, E. Ben-Jacob and S. Efrati (2015). "Hyperbaric oxygen may induce angiogenesis in patients suffering from prolonged post-concussion syndrome due to traumatic brain injury." Restorative neurology and neuroscience **33**(6): 943-951.

van der Kolk, B. A., S. Roth, D. Pelcovitz, S. Sunday and J. Spinazzola (2005). "Disorders of extreme stress: The empirical foundation of a complex adaptation to trauma." J Trauma Stress **18**(5): 389-399.

Vera, P., M. Zilbovicius, H. Chabriat, P. Amarenco, J. Kerdraon, J. F. Menard, G. Amarenco, M. G. Bousser, G. Rancurel and Y. Samson (1996). "Post-stroke changes in cortical 5-HT₂ serotonergic receptors." J Nucl Med **37**(12): 1976-1981.

Vilela, D. S., P. R. Lazarini and C. F. Da Silva (2008). "Effects of hyperbaric oxygen therapy on facial nerve regeneration." Acta Otolaryngol **128**(9): 1048-1052.

Yamauchi, H., T. Kudoh, Y. Kishibe, J. Iwasaki and S. Kagawa (2005). "Selective neuronal damage and borderzone infarction in carotid artery occlusive disease: a 11C-flumazenil PET study." J Nucl Med **46**(12): 1973-1979.

Yang, Y. J., X. L. Wang, X. H. Yu, X. Wang, M. Xie and C. T. Liu (2008). "Hyperbaric oxygen induces endogenous neural stem cells to proliferate and differentiate in hypoxic-ischemic brain damage in neonatal rats." Undersea Hyperb Med **35**(2): 113-129.

Yavne, Y., D. Amital, A. Watad, S. Tiosano and H. Amital (2018). "A systematic review of precipitating physical and psychological traumatic events in the development of fibromyalgia." Semin Arthritis Rheum.

Yildiz, S., M. Z. Kiralp, A. Akin, I. Keskin, H. Ay, H. Dursun and M. Cimsit (2004). "A new treatment modality for fibromyalgia syndrome: hyperbaric oxygen therapy." J Int Med Res **32**(3): 263-267.

Yildiz, S., M. Z. Kiralp, A. Akin, I. Keskin, H. Ay, H. Dursun and M. Cimsit (2004). "A new treatment modality for fibromyalgia syndrome: hyperbaric oxygen therapy." The Journal of international medical research **32**(3): 263-267.

Yin, W., A. E. Badr, G. Mychaskiw and J. H. Zhang (2002). "Down regulation of COX-2 is involved in hyperbaric oxygen treatment in a rat transient focal cerebral ischemia model." Brain Res **926**(1-2): 165-171.

YOSHIHIRO MASAKI, T. N., YOSHITAKA IKEJIRI, HIROMASA TOKUNAGA, NAOKI HATTA, TAKESHI UEMA, HIROAKI KAZUI, TALANT K. DORONBEKOV, ATSUSHI, OGINO, NORIKO MIYOSHI, HISASHI TANII, TOSHIHISA TANAKA, NAOHIKO OKU (2006). "Neural substrates of emotionally valenced episodic memory: A PET study using film stimuli." Psychiatry and Clinical Neurosciences.

Yunus, M. B. (2007). "Fibromyalgia and overlapping disorders: the unifying concept of central sensitivity syndromes." Seminars in arthritis and rheumatism **36**(6): 339-356.

Yunus, M. B. (2007). "Role of central sensitization in symptoms beyond muscle pain, and the evaluation of a patient with widespread pain." Best practice & research. Clinical rheumatology **21**(3): 481-497.

In review

Figure 1.TIF




Figure 2.TIF




Figure 3.TIF

HBOT



Increased tissue oxygenation



Anaerobic-> Aerobic metabolism



Brain regeneration processes



Increased cerebral blood flow



Increased cortical activity/metabolism



Suppressed memory recovery

In review